



Causes of Revision of Total Knee Arthroplasties in a Tertiary Hospital in Brazil

Causas de revisão de artroplastias totais do joelho em um hospital terciário no Brasil

Alan de Paula Mozella¹ Hugo Alexandre de Araújo Barros Cobra¹ Sandra Tie Nishibe Minamoto¹
Rodrigo Salim² Ana Carolina Leal³

¹Department of Orthopedics and Traumatology, Knee Surgery Center, Instituto Nacional de Traumatologia e Ortopedia (INTO), Rio de Janeiro, RJ, Brazil

²Department of Orthopedics and Anesthesiology, Faculdade de Medicina de Ribeirão Preto, Universidade de São Paulo (USP), São Paulo, SP, Brazil

³Division of Teaching and Research, Instituto Nacional de Traumatologia e Ortopedia (INTO), Rio de Janeiro, RJ, Brazil

Address for correspondence Alan de Paula Mozella, Instituto Nacional de Traumatologia e Ortopedia, (INTO), Rio de Janeiro, RJ, Brazil (e-mail: apmozella@terra.com.br).

Rev Bras Ortop 2024;59(5):e696–e701.

Abstract

Objective To identify the causes of revision of total knee arthroplasty in a referral center in Brazil.

Methods This is a case series, with 80 patients undergoing revision surgery for total knee arthroplasty (RTKA) at a referral center for knee surgery, between August 2019 and November 2021, with a mean age of 69.6 years. Of these patients, 60.23% were female and 39.77% were male. The average body mass index (BMI) was 30.23 kg/m². The causes of TKA failure were defined as: periprosthetic infection according to the 2018 International Consensus Meeting criteria, ligament instability, range of motion limitation, periprosthetic fracture, malalignment, aseptic loosening, pain due to non-replacement of the patellar cartilage, polyethylene wear, fracture of implants, insufficiency of the extensor mechanism.

Results Periprosthetic joint infection (PJI) was the main cause of revision total knee arthroplasty (TKA), corresponding to 47.73% of cases. Aseptic loosening of one or more components represented the second most frequent reason for TKA failure, accounting for 35.23% of revisions. Range of motion limitation represented the third most frequent cause, accounting for 5.68% of surgeries. Instability was the fourth most frequent reason for RTKA, occurring in 4.55% of patients. The other causes of revision were: periprosthetic fracture (3.41%), failure due to rupture of the extensor mechanism (2.27%), and pain attributed to non-replacement of the patellar cartilage (1.14%).

Keywords

- ▶ arthroplasty
- ▶ arthroplasty, replacement, knee
- ▶ infections
- ▶ knee joint

Work carried out at the Instituto Nacional de Traumatologia e Ortopedia (INTO), Rio de Janeiro, RJ, Brazil.

received
April 19, 2022
accepted
August 17, 2022

DOI <https://doi.org/10.1055/s-0042-1757304>.
ISSN 0102-3616.

© 2024. The Author(s).

This is an open access article published by Thieme under the terms of the Creative Commons Attribution 4.0 International License, permitting copying and reproduction so long as the original work is given appropriate credit (<https://creativecommons.org/licenses/by/4.0/>).

Thieme Revinter Publicações Ltda., Rua do Matoso 170, Rio de Janeiro, RJ, CEP 20270-135, Brazil

Conclusions Periprosthetic joint infection was the most frequent cause of TKA revision in our series. Other reasons for TKA failures were, in descending order: aseptic loosening, limited range of motion, and instability.

Resumo

Objetivo Identificar as causas de revisão de artroplastia total do joelho em um centro de referência no Brasil.

Métodos Trata-se de uma série de casos, com 80 pacientes submetidos à cirurgia de revisão de artroplastia total do joelho (RATJ) em um centro de referência em cirurgia do joelho, no período entre agosto de 2019 e novembro de 2021, com média de idade de 69,6 anos. Desses pacientes, 60,23% eram do sexo feminino e 39,77% do sexo masculino. O índice de massa corporal (IMC) médio foi de 30,23 kg/m². As causas de falha da ATJ foram definidas como: infecção periprotética conforme critérios do 2018 *Internacional Consensus Meeting*, instabilidade ligamentar, limitação de arco de movimento, fratura periprotética, mal alinhamento, soltura asséptica, dor devido à não substituição da cartilagem patelar, desgaste do polietileno, fratura dos implantes, insuficiência do mecanismo extensor.

Resultados A infecção articular periprotética (IAP) foi a principal causa de revisão de artroplastia total do joelho (ATJ), correspondendo a 47,73% dos casos. A soltura asséptica de um ou mais componentes representou o segundo motivo mais frequente para causa de falha de ATJ, acarretando 35,23% das revisões. Limitação do arco de movimento representou a terceira mais frequente causa, sendo responsável por 5,68% das cirurgias. Instabilidade foi a 4ª razão mais frequente para RATJ, ocorrendo em 4,55% dos pacientes. As demais causas de revisão foram: fratura periprotética (3,41%), insuficiência por ruptura do mecanismo extensor (2,27%), dor atribuída à não substituição da cartilagem patelar (1,14%).

Conclusões A infecção articular periprotética foi a causa mais frequente de revisão de ATJ em nossa série. Outras razões de falhas de ATJ foram, em ordem decrescente: soltura asséptica, limitação do arco de movimento e instabilidade.

Palavras-chave

- articulação do joelho
- artroplastia
- artroplastia do joelho
- infecções

Introduction

Epidemiological data show that between 6 and 12% of total knee arthroplasties (TKA) evolve to failure, requiring revision surgery in the first 10 years.¹ Other recent studies have shown an increase in the number of revision surgeries of TKA (RTKA) performed in recent years.²⁻⁵ There was a 144% increase in the number of RTKA performed in Germany between 2004 and 2014.⁶ It is estimated that, for the year 2050, in the same country, there will be an increase of about 90% in the number of revision surgeries of knee arthroplasties.⁴ In the United States, the increase in the number of such procedures was of 39% between 2006 and 2010.⁷ And future estimates point to a continuous increase in the number of these surgeries by the year 2030.⁸

The etiology for increasing the number of RTKAs is, in fact, multifactorial, being influenced both by factors related to technical issues and evolution of implants and instruments, as well as by factors related to patients.^{9,10} An example of this, the wear of the polyethylene component was a frequent cause of revision in the past; however, changes in the sterilization process of these components culminated in a significant reduction in the number of revisions for this

reason.¹¹⁻¹³ At the same time, the expansion of primary prosthesis indications, especially for younger patients, and the higher incidence of obesity in the world population may, at least partially, also justify potential changes in the failure patterns of modern arthroplasties.^{9,14,15} Meehan et al.¹⁴ demonstrated that the risk of septic failure was 1.81 times higher in patients under 50 years of age when compared to patients older than 65 years. Similarly, among younger patients, the risk of revision for aseptic failure was 4.7 times higher.¹⁴

Knee prosthesis revision surgeries are procedures of high complexity, performed in a limited number of hospitals and, consequently, that require greater expenditure of technical and economic resources.^{16,17} Compliance with technical principles is essential to obtain satisfactory results in this type of surgery.¹⁸ Thus, preoperative planning, failure mode identification and adequate treatment of bone defects are of fundamental importance for obtaining satisfactory and long-lasting clinical results in this type of surgery.^{18,19}

In this context, the identification of the cause of implant failure is significantly important for planning and obtaining satisfactory results.²⁰ Several series have demonstrated

changes in the causes of TKA failure over the years, possibly related to evolutions in implant design, as well as instrumental improvements.^{11,12} Thus, this study aims to identify the causes of RTKA in a reference center in Brazil.

Materials and Methods

This is a series of cases of patients undergoing RTKA at the Center for Specialized Attention in Knee Surgery of Instituto Nacional de Traumatologia e Ortopedia (INTO) Jamil Haddad, in the period between August and November 2019 (CAAE-20309419.0.0000.5273.). There was no age limit, no restriction on the gender of the participants.

During the study period, 88 arthroplasty revision surgeries were performed. One patient underwent revision of the unicompartmental implant of the knee and was therefore excluded. Seven surgeries were excluded from the analysis because it was the second revision surgery. Thus, after the exclusion criteria, 80 surgeries (80 patients) were analyzed. The mean age of the patients in the procedures was 69.6 years (standard deviation (SD) = 9.85 years; variation = 29–87 years). Of these patients, 60.23% were female and 39.77% were male. The mean body mass index (BMI) was 30.23 kg/m² (SD = 5.92; variation = 18.20–42.46 kg/m²).

Arthroplasty review was defined as reoperation after performing primary TKA in which it was necessary to add or replace one or more prosthetic components. Thus, patients submitted to reoperations, however without the addition or replacement of one or more primary prosthetic components, were excluded from the analysis. Patients submitted to unicompartmental knee implant (UKI) revision surgery were also excluded from the analysis, as well as patients submitted to the second revision surgery.

The following data were collected from medical records: age, gender, and BMI. Laboratory tests with analysis of hemossedimentation velocity (HSV), C-reactive protein (CRP), and D-dimer were performed in all patients on the day before surgery, as well as range of motion measurement using a goniometer by the orthopedist in training in the knee surgery improvement course at the institution. Radiographs of the knee at anteroposterior incidences with bipodal, profile, and axial support in 30 degrees were performed on the day before surgery. During surgery, synovial fluid was collected for the following analyses: total leukocyte count, polymorphonuclear percentage, leukocyte esterase, and culture in a blood culture vial. Six samples of bone tissue were also collected for extended tissue culture for 14 days and 2 periprosthetic membrane samples for histopathological examination.

In the surgical report, the following information were analyzed: surgical access performed, implant system used, Anderson Orthopaedic Research Institute (AORI) classification of bone loss, use of trabeculated metal cones and/or tissue bank grafts for the treatment of defects.

Regarding the surgical access used, the surgery was performed by conventional medial parapatellar access in 70 surgeries (87.5%). Extended access was performed in 10 surgeries (12.5%), with tibia anterior tuberosity (TAT) osteot-

Table 1 Distribution of bone defects

Classification of defects	Tibia n (%)	Femur n (%)
Flawless	15 (20%)	16 (21.3%)
1	20 (26.7%)	15 (20%)
2a	19 (25.3%)	15 (20%)
2b	9 (12%)	8 (10.7%)
3	12 (16%)	21 (28%)

Source: Research data (INTO).

omy in 7 cases (8.75%), quadriceps turndown in two cases (2.5%), and clipping of the thigh reskin (snip) in one patient (1.25%).

Implants with two different systems—Legacy Constrained Condylar Knee (Zimmer Biomet, Warsaw, IN, USA) or Legion Revision Knee (Smith & Nephew plc, London, UK)—were used in 40 surgeries (50%). In 24 surgeries (30%), constrictor implants (Rotating Hinge Knee – Zimmer Biomet) were used. Spacers were implanted in 14 patients (17.5%). In 2 patients (2.5%), only patellar implants were used.

Medical documentation of bone defects was incomplete in five surgeries. Thus, when analyzing 75 patients, the presence of bone defects was identified in 70% of the patients. The distribution and classification of bone failures are shown in ►Table 1. Tantalum metaphysary cones were necessary for the treatment of bone defects in 21 patients (26.25%). Of these, tissue bank grafting was also used in nine surgeries. In all, the homologous graft was used in 13 patients (16.25%).

The cause of RTKA was defined by the surgeon responsible for the procedure based on the following criteria: all patients were submitted to evaluation according to the criteria of the 2018 International Consensus Meeting (ICM 2018).^{21,22} The diagnosis of periprosthetic infection was made when at least one of the major criteria was present, that is, the presence of a fistula communicating with the joint or presence of the same pathogen in two or more cultures of periimplant tissues. The diagnosis of periprosthetic infection using the lower criteria was confirmed when the score of the sum of the criteria was greater than or equal to six when the parameters of the tests recommended by the ICM 2018 were observed. In cases defined with aseptic failures, the surgeon in charge, after analysis of the physical examination, imaging and laboratory examinations defined the reason for failure between the following possible: ligament instability (when presence of dislocation or subluxation of prosthetic components, or when pathological ligament opening greater than 5 mm during physical examination, or ligament laxity greater than 5 mm in varus or varus stress tests during physical examination or ap radiographs of the knee with load), limitation of range of motion (when less than 50 degrees associated with physical disability declared by the patient), periprosthetic fracture, poor alignment, aseptic loosening, pain due to non-replacement of patellar cartilage, polyethylene wear, fracture of implants, or insufficiency of the extensor mechanism.




Fig. 1 Distribution of the causes of failure of total knee arthroplasty. Causa da Cirurgia = Reason for surgery, Infecção = Infection, Soltura Asséptica = Aseptic loosening, Instabilidade = Instability, Dor em patela não substituída = Pain on non-replaced patella, Fratura peri-protética = Peri-prosthetic fracture, Lesão do mecanismo extensor = Extensive mechanism injury, Limitação da ADM (artrofibrose) = ROM limitation (arthrofibrosis)

Results

Periprosthetic joint infection (PJI) was the main cause of RTKA, corresponding to 47.73% of cases. The aseptic loosening of one or more components represented the second most frequent cause of failure, resulting in 35.23% of revisions. Limitation of the range of motion represented the 3rd most frequent cause, accounting for 5.68% of surgeries. Instability was the 4th most frequent cause of failure, occurring in 4.55% of patients. The other causes of revision were: periprosthetic fracture (3.41%), insufficiency due to rupture of the extensor mechanism (2.27%), pain attributed to non-replacement of patellar cartilage (1.14%) (→ Fig. 1).

Discussion

The main finding of our study was to identify that the main cause of failure in our series of primary TKAs was periprosthetic infection. Other reasons that led to the need for revision were, in decreasing order, aseptic loosening, limitation of the range of motion, and joint instability. Similarly, several recent studies have demonstrated periprosthetic infection as the most frequent mechanism of failure of primary prostheses, reaching 20 to 37.7% of surgeries.^{2,23–25} In the study by Evangelopoulos et al.,²⁴ periprosthetic infection was the main mechanism of failure with an incidence of 26.3%. However, when analyzing only patients submitted to the second revision surgery, the authors identified that the incidence of periprosthetic infection remained the most frequent reason for TKA failure with approximately 50% of cases. We believe that this high incidence of revisions due to septic failure observed in our study may have been influenced, at least partially, by the fact that the surgeries were performed in the COVID-19 pandemic period. In this scenario, in several months, only emergency surgeries were performed, such as acute periprosthetic infections and, consequently, subsequent increase in the demand for reimplantation after treatment of infection in two times.

Aseptic loosening represented the 2nd most frequent reason for implant failure in our series, occurring in approxi-

mately 35% of cases. Similarly, Bozic et al.⁷ demonstrated that periprosthetic infection and aseptic loosening were the two most frequent reasons for TKA revision. When analyzing more than 60,000 primary surgeries, the authors identified septic failure in 25.1% of patients and aseptic loosening in 16.7% of cases.⁷ Our results are in agreement with those of Koh et al.²⁶ and Evangelopoulos et al.,²⁴ who demonstrated aseptic loosening as the 2nd main cause of revision, affecting, respectively, 32.7% and 25% of the failures. Siqueira et al.,¹ when analyzing the arthroplasty registry of five countries, identified aseptic loosening as the first cause of revision, affecting 30% of surgeries. Several other studies have also reported aseptic loosening as the main mechanism of TKA failure with incidence ranging from 24 to 44%.^{10,27,28} We identified that the incidence of aseptic loosening in our series is in accordance with the literature, although it is not the main mode of failure of our implants.

The limitation of the range of motion represented, in our series, the third most common indication for RTKA, with approximately 6% of surgeries. Similarly, in the study by Le et al., arthrofibrosis was the third most frequent indication for revision, although the incidence presented corresponds to approximately three times that observed in our study. Pietrzak et al.¹¹ studied only the aseptic causes of revision and identified joint stiffness as the 2nd leading cause of revision with an incidence of 27.5%. Meanwhile, Le²⁹ and Koh et al.²⁶ showed an incidence that only 2.5% of patients required TKA revision due to functional limitation imposed by movement arc restriction. Such differences can be explained, at least partially, by the various definitions of joint stiffness, as well as being influenced by the functional demand and expectations of patients.^{30–32} We should also observe that, in our series, we included only patients submitted to revision of prosthetic components; thus, patients submitted to manipulation under narcosis after TKA or even reoperation with joint release were not analyzed, which could have modified the incidence if we analyzed reoperations after knee prosthesis. In a study conducted in Brazil evaluating short-term complications (up to 1 year), the authors identified that joint stiffness was the most frequent complication affecting 7.5% of patients undergoing primary knee arthroplasty.³³

In our study, component instability represented the 4th cause of RTKA, with approximately 5% of surgeries. Other studies have demonstrated an incidence of instability ranging from 18.7 to 30%, constituting the 2nd or 3rd most frequent reason for TKA review.^{10,28,29} However, in the works of Kasahara et al.²⁷ and Koh et al.,²⁶ instability was identified in 9.3% and 6.5%, respectively. Post-TKA instability is usually indicated as a cause of early failure, that is, in implants with less than 2 years of evolution.^{6,11} Thus, the reduction in the number of elective primary surgeries performed during the study period, due to the restrictions imposed by the COVID-19 pandemic, may justify, at least partially, the limited number of revisions due to post-TKA instability. Another potential explanation can be attributed to the waiting time for this type of surgery in the Brazilian public health service. Thus, any

initial cases of instability can potentially evolve with loosening of the implants, thus hindering the primary diagnosis of component instability.

Despite the importance of our series reporting the distribution of the different causes of RTKA in the national scenario, our study has significant limitations. We believe that restrictions on elective surgeries imposed during part of the COVID-19 pandemic period may influence the incidence of some modes of failure, such as suspension of elective aseptic revisions in certain periods. Another important limitation refers to the retrospective nature of our analysis. Thus, several patients submitted to revision surgery had been operated on primarily in other hospitals. Thus, we had no control over the evolution time from the implantation of the components to the occurrence of failure requiring revision surgery, nor did we perform analyses of the different systems of primary prostheses implanted. Schroer et al.¹⁰ indicate that 35.3% of knee arthroplasties fail in the first 2 years after implantation, being more often related to factors influenced by the surgeon than by the performance of implants. However, we understand that such information is of high relevance for the planning of the Brazilian health system in view of the increase in the number of cases of RTKA, given, the impacts regarding the need for hospital beds, financial resources, and technical training of the teams. Similarly, we believe that the high number of revision surgeries performed due to septic failure observed in our series alerts us to the need to expand the measures to prevent these serious complications.

Therefore, we believe that the adoption of preventive measures of postoperative infection, such as: optimization of preoperative clinical conditions, adoption of risk stratification scores and investigation of patients colonized by *Methicillin-resistant Staphylococcus aureus* may impact on the reduction of the number of RTKA surgeries performed in our hospital.

Conclusions

We identified periprosthetic joint infection as the most frequent cause of RTKA. The other TKA failure mechanisms were, in decreasing order: aseptic loosening, limitation of the range of motion, and instability. Periprosthetic fracture and extensor mechanism insufficiency were less frequent causes of primary prosthesis revision in our series.

Financial Support

The authors declare that there was no financial support from public, commercial, or non-profit sources.

Conflict of Interests

The authors declare that there is no conflict of interests.

References

- 1 Siqueira MBP, Klika AK, Higuera CA, Barsoum WK. Modes of failure of total knee arthroplasty: registries and realities. *J Knee Surg* 2015;28(02):127–138
- 2 Mathis DT, Lohrer L, Amsler F, Hirschmann MT. Reasons for failure in primary total knee arthroplasty - An analysis of prospectively collected registry data. *J Orthop* 2020;23:60–66
- 3 Geary MB, Macknet DM, Ransone MP, Odum SD, Springer BD. Why Do Revision Total Knee Arthroplasties Fail? A Single-Center Review of 1632 Revision Total Knees Comparing Historic and Modern Cohorts. *J Arthroplasty* 2020;35(10):2938–2943
- 4 Rupp M, Walter N, Lau E, Worlicek M, Kurtz SM, Alt V. Recent trends in revision knee arthroplasty in Germany. *Sci Rep* 2021;11(01):15479
- 5 Klug A, Gramlich Y, Rudert M, et al. The projected volume of primary and revision total knee arthroplasty will place an immense burden on future health care systems over the next 30 years. *Knee Surg Sports Traumatol Arthrosc* 2021;29(10):3287–3298
- 6 Boelch SP, Jakuscheit A, Doerries S, et al. Periprosthetic infection is the major indication for TKA revision - experiences from a university referral arthroplasty center. *BMC Musculoskelet Disord* 2018;19(01):395
- 7 Bozic KJ, Kurtz SM, Lau E, et al. The epidemiology of revision total knee arthroplasty in the United States. *Clin Orthop Relat Res* 2010;468(01):45–51
- 8 Kurtz S, Ong K, Lau E, Mowat F, Halpern M. Projections of primary and revision hip and knee arthroplasty in the United States from 2005 to 2030. *J Bone Joint Surg Am* 2007;89(04):780–785
- 9 Kamath AF, Ong KL, Lau E, et al. Quantifying the Burden of Revision Total Joint Arthroplasty for Periprosthetic Infection. *J Arthroplasty* 2015;30(09):1492–1497
- 10 Schroer WC, Berend KR, Lombardi AV, et al. Why are total knees failing today? Etiology of total knee revision in 2010 and 2011. *J Arthroplasty* 2013;28(8, Suppl):116–119
- 11 Pietrzak J, Common H, Migaud H, Pasquier G, Girard J, Putman S. Have the frequency of and reasons for revision total knee arthroplasty changed since 2000? Comparison of two cohorts from the same hospital: 255 cases (2013–2016) and 68 cases (1991–1998). *Orthop Traumatol Surg Res* 2019;105(04):639–645
- 12 Sharkey PF, Lichstein PM, Shen C, Tokarski AT, Parvizi J. Why are total knee arthroplasties failing today—has anything changed after 10 years? *J Arthroplasty* 2014;29(09):1774–1778
- 13 Thiele K, Perka C, Matziolis G, Mayr HO, Sostheim M, Hube R. Current failure mechanisms after knee arthroplasty have changed: polyethylene wear is less common in revision surgery. *J Bone Joint Surg Am* 2015;97(09):715–720
- 14 Meehan JP, Danielsen B, Kim SH, Jamali AA, White RH. Younger age is associated with a higher risk of early periprosthetic joint infection and aseptic mechanical failure after total knee arthroplasty. *J Bone Joint Surg Am* 2014;96(07):529–535
- 15 Jämsen E, Nevalainen P, Eskelinen A, Huotari K, Kalliovaara J, Moilanen T. Obesity, diabetes, and preoperative hyperglycemia as predictors of periprosthetic joint infection: a single-center analysis of 7181 primary hip and knee replacements for osteoarthritis. *J Bone Joint Surg Am* 2012;94(14):e101
- 16 Morcos MW, Kooner P, Marsh J, Howard J, Lanting B, Vasarhelyi E. The economic impact of periprosthetic infection in total knee arthroplasty. *Can J Surg* 2021;64(02):E144–E148
- 17 Boddapati V, Fu MC, Mayman DJ, Su EP, Sculco PK, McLawhorn AS. Revision Total Knee Arthroplasty for Periprosthetic Joint Infection Is Associated With Increased Postoperative Morbidity and Mortality Relative to Noninfectious Revisions. *J Arthroplasty* 2018;33(02):521–526
- 18 Mozella AP, Cobra HAAB. Bone Defects in Revision Total Knee Arthroplasty. *Rev Bras Ortop* 2021;56(02):138–146
- 19 Sheth NP, Bonadio MB, Demange MK. Bone Loss in Revision Total Knee Arthroplasty: Evaluation and Management. *J Am Acad Orthop Surg* 2017;25(05):348–357
- 20 Pitta M, Esposito CI, Li Z, Lee YY, Wright TM, Padgett DE. Failure After Modern Total Knee Arthroplasty: A Prospective Study of 18,065 Knees. *J Arthroplasty* 2018;33(02):407–414

- 21 Schwarz EM, Parvizi J, Gehrke T, et al. 2018 International Consensus Meeting on Musculoskeletal Infection: Research Priorities from the General Assembly Questions. *J Orthop Res* 2019;37(05):997–1006
- 22 Parvizi J, Tan TL, Goswami K, et al. The 2018 Definition of Periprosthetic Hip and Knee Infection: An Evidence-Based and Validated Criteria. *J Arthroplasty* 2018;33(05):1309–1314.e2
- 23 Delanois RE, Mistry JB, Gwam CU, Mohamed NS, Choksi US, Mont MA. Current Epidemiology of Revision Total Knee Arthroplasty in the United States. *J Arthroplasty* 2017;32(09):2663–2668
- 24 Evangelopoulos DS, Ahmad SS, Krismer AM, et al. Periprosthetic Infection: Major Cause of Early Failure of Primary and Revision Total Knee Arthroplasty. *J Knee Surg* 2019;32(10):941–946
- 25 Meyer JA, Zhu M, Cavadino A, Coleman B, Munro JT, Young SW. Infection and periprosthetic fracture are the leading causes of failure after aseptic revision total knee arthroplasty. *Arch Orthop Trauma Surg* 2021;141(08):1373–1383
- 26 Koh IJ, Cho WS, Choi NY, Kim TKKleos Korea Research Group. Causes, risk factors, and trends in failures after TKA in Korea over the past 5 years: a multicenter study. *Clin Orthop Relat Res* 2014;472(01):316–326
- 27 Kasahara Y, Majima T, Kimura S, Nishiike O, Uchida J. What are the causes of revision total knee arthroplasty in Japan? *Clin Orthop Relat Res* 2013;471(05):1533–1538
- 28 Mulhall KJ, Ghomrawi HM, Scully S, Callaghan JJ, Saleh KJ. Current etiologies and modes of failure in total knee arthroplasty revision. *Clin Orthop Relat Res* 2006;446(446):45–50
- 29 Le DH, Goodman SB, Maloney WJ, Huddleston JI. Current modes of failure in TKA: infection, instability, and stiffness predominate. *Clin Orthop Relat Res* 2014;472(07):2197–2200
- 30 Rutherford RW, Jennings JM, Levy DL, Parisi TJ, Martin JR, Dennis DA. Revision Total Knee Arthroplasty for Arthrofibrosis. *J Arthroplasty* 2018;33(7S):S177–S181
- 31 Cheuy VA, Foran JRH, Paxton RJ, Bade MJ, Zeni JA, Stevens-Lapsley JE. Arthrofibrosis Associated With Total Knee Arthroplasty. *J Arthroplasty* 2017;32(08):2604–2611
- 32 Thompson R, Novikov D, Cizmici Z, et al. Arthrofibrosis After Total Knee Arthroplasty: Pathophysiology, Diagnosis, and Management. *Orthop Clin North Am* 2019;50(03):269–279
- 33 Carvalho LH Junior, Castro CAC, Gonçalves MJB, Rodrigues LCM, Lopes FL, Cunha FVP. Complicações de curto prazo da artroplastia total do joelho: avaliação de 120 casos. *Rev Bras Ortop* 2006;41(05):162–166